NAVAIR 50-1C-7
AWS/TR-89/007

JOINT U.S. NAVY/U.S. AIR FORCE CLIMATIC STUDY OF THE UPPER ATMOSPHERE

VOLUME 7 - JULY

JULY, 1989

AD-A227 126



PREPARED BY
NAVAL OCEANOGRAPHY COMMAND DETACHMENT
ASHEVILLE, N.C.

DTIC
ELECTE
OCT 04 1990
S E D

PREPARED UNDER THE AUTHORITY OF
COMMANDER, NAVAL OCEANOGRAPHY COMMAND
STENNIS SPACE CENTER, MS 39529-5000

BEST
AVAILABLE COPY

90 10 03 095

0850LP0158000

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

NAVAIR 50-1C-7
AWS/TR-89/007

JOINT U.S. NAVY/U.S. AIR FORCE CLIMATIC STUDY OF THE UPPER ATMOSPHERE

VOLUME 7 - JULY

JULY, 1989



PREPARED BY
NAVAL OCEANOGRAPHY COMMAND DETACHMENT
ASHEVILLE, N.C.

DTIC
ELECTE
OCT 04 1990
S E D

PREPARED UNDER THE AUTHORITY OF
COMMANDER, NAVAL OCEANOGRAPHY COMMAND
STENNIS SPACE CENTER, MS 39529-5000

90 1 005

0850LP0158000

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified	1b RESTRICTIVE MARKINGS
2a SECURITY CLASSIFICATION AUTHORITY	3 DISTRIBUTION / AVAILABILITY OF REPORT Public Release/Distribution Unlimited
2b DECLASSIFICATION / DOWNGRADING SCHEDULE	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)	5 MONITORING ORGANIZATION REPORT NUMBER(S) NAVAIR 50-1C-7 S/N 0850-LP-015-8000, AWS/TR-89/007
6a NAME OF PERFORMING ORGANIZATION National Climatic Data Center Global Analysis Branch 6b OFFICE SYMBOL (If applicable) E/CC22	7a NAME OF MONITORING ORGANIZATION Naval Oceanography Command Detachment Asheville
6c ADDRESS (City, State, and ZIP Code) Federal Building Asheville, NC 28801-2696	7b ADDRESS (City, State, and ZIP Code) Federal Building Asheville, NC 28801-2696
8a NAME OF FUNDING / SPONSORING ORGANIZATION Commander, Naval Oceanography Command Headquarters, Air Weather Service	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER
8c ADDRESS (City, State, and ZIP Code) Stennis Space Center, MS 39529-5000 Scott AFB, IL 62225-5008	10 SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO / PROJECT NO / TASK NO / WORK UNIT ACCESSION NO

11 TITLE (Include Security Classification)
Joint U.S. Navy/U.S. Air Force Climatic Study of the Upper Atmosphere Volume 7-July

12 PERSONAL AUTHOR(S) NCDC - Michael J. Changery, Claude N. Williams
NAVOCEANCOMDET - Michael L. Dickenson, Brian L. Wallace

13a TYPE OF REPORT	13b TIME COVERED	14 DATE OF REPORT (Year, Month, Day)	15 PAGE COUNT
Final	FROM _____ TO _____	July 1989	236

16 SUPPLEMENTARY NOTATION

17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
FIELD	GROUP	SUB-GROUP	

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

This study of the upper atmosphere is based on 1980-85 twice daily gridded analysis produced by the European Centre for Medium Range Weather Forecasts. Included are global analyses of (1) Mean Temperature/Standard Deviation, (2) Mean Geopotential Height/Standard Deviation, (3) Mean Density/Standard Deviation, (4) Height and Vector Standard Deviation. All for 13 pressure levels - 1000, 850, 700, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30 mb. In addition, analyses of (5) Mean Dew Point/Standard Deviation - levels 1000 through 300 mb, (6) jet stream (mean scalar speed) - levels 500 through 30 mb. Also included are global 5 degree grid point wind roses for the 13 pressure levels.

20 DISTRIBUTION / AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS	21 ABSTRACT SECURITY CLASSIFICATION Unclassified
22a NAME OF RESPONSIBLE INDIVIDUAL Brian L. Wallace	22b TELEPHONE (Include Area Code) (704) 252-7865 22c OFFICE SYMBOL

DD FORM 1473, 84 MAR 83 APR edition may be used until exhausted
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

TABLE OF CONTENTS

	PAGE
INTRODUCTION	iv
REFERENCES	vii
ELEMENTS	
PRESSURE-HEIGHT	1-27
WIND ROSE	29-107
JET STREAM	109-129
TEMPERATURE	131-157
DEW POINT	159-171
DENSITY	173-199
HEIGHT/WIND STANDARD DEVIATION	201-227

Accession For
NTIS GRA&I
DTIC TAB
Unannounced
Justification
By
Distribution/
Availability Codes
Dist | Avail and/or Special
A-1

The Joint U.S. Navy/U.S. Air Force Climatic Study of the Upper Atmosphere was prepared by the Officer in Charge, Naval Oceanography Command Detachment, Asheville, North Carolina under the authority of Commander, Naval Oceanography Command. Additional funding was provided by the Air Weather Service as a result of Tri-Services Climatology initiatives. The work was performed at the National Climatic Data Center (NCDC). Specific acknowledgement of the NCDC staff is made to Mr. M.J. Changery, project leader; Mr. C.N. Williams, Jr. for data processing and software development; and Messrs. M.G. Burgin and D.A. McKittrick for drafting skills. Special acknowledgement is made to the European Centre for Medium-range Weather Forecasts for providing the basic gridded analyses.

INTRODUCTION

During the past decade, improvements in the collection and assimilation of data required for more accurate representations of the atmosphere have resulted in data sets useful for developing a more definitive climatology of the global atmosphere. Such a climatology has uses in aircraft operations and planning, indirect assessments of atmospheric transport as well as a standard state from which atmospheric anomalies can be analyzed.

Prior climatologies, U.S. Navy (1959), U.S. Navy (1966), Naval Weather Service Command (1969), and Naval Weather Service Command (1970), were produced from individual station data with varying periods of record, and the resulting summarized data were analyzed. A serious deficiency was the lack of reporting locations in the major ocean basins. Analyses over the oceans were derived by extrapolating from known analyses over coastal regions as well as the few island or ocean vessels available. An additional complication was the manually intensive effort required to ensure horizontal and vertical consistency of the data.

With the advent, in the 1970s, of more powerful computers and data collection and assimilation systems, the initial analyses used for input into forecast models had a three-fold advantage over the station analyses utilized in the prior climatologies. First, the data assimilation system utilized a greater variety of information for production of an analysis. The normal array of land-based upper air reporting stations was supplemented by ship-based reporting stations, cloud reports, pilot reports and, most importantly, satellite-derived temperature, moisture and wind data. Consequent analyses more accurately represented the state of the atmosphere at a given observation time. Second, the assimilation system quality-controlled all incoming data and ensured the horizontal and vertical consistency of the resulting analyses. Finally, through the computer-based system, global data were available and archived in grid-point form.

A number of analysis sets produced by various national and international meteorological services were investigated. It is recognized that improvements to the data assimilation and analysis systems occurred within any analysis set produced, and that current analyses more accurately reflect the atmosphere's state than do the earlier analyses. It is also recognized that specific parameter or geographic-based deficiencies exist in all analysis sets. However, the intent of this upper-air climatology effort is the production of analyses to serve the needs of the operational meteorologist. A climatology derived from global analyses achieves this goal. Based on known capabilities and technical reviews of the various systems, as well as recommendations from the professional numerical modeling community, the analyses produced by the European Centre for Medium-range Forecasts were selected for processing.

ECMWF DATA

The European Centre for Medium-range Weather Forecasts (ECMWF) is an international organization established in 1973 and supported by 17 member states. It is responsible for providing global forecasts to the European community. Their data assimilation system consists of multivariate optimal interpolation analysis allowing the incorporation of a variety of observations with differing error characteristics and spatial distributions. A relatively comprehensive coverage of global data is ensured through the data collection schedule. A unique feature of the ECMWF system is the method of grid point analysis. Rather than analyzing individual grid points, varying sized boxes (depending on data density) are created containing groups of grid points. Grid point analysis uses data from within the box as well as adjacent boxes, thereby assuring a consistent analysis between all the grid points.

The system also includes internal quality control which examines the climatological reasonability of incoming data as well as the internal consistency of the data.

In addition, the system utilizes a model initialization process which ensures that harmful gravity waves, caused by imbalances in the analysis, with the potential to create problems in subsequent forecast fields, are suppressed. Through the initialization process, the atmosphere's mass and wind fields are adjusted so that only a portion of the gravity wave balanced by dynamic and physical processes is retained. Further information on the ECMWF system is available in Lorenc (1981), Shaw, et al. (1984), Lonnberg, et al. (1986), and ECMWF (1988).

The resulting initialized analyses are vertically interpolated to these 13 standard pressure levels: 1000, 850, 700, 500, 400, 300, 250, 200, 150, 100, 70, 50, and 30 mb, and include the geopotential height, temperature, and wind for all levels with moisture included for the 1000 through 300 mb levels.

Six years (1980-1985) of individual analysis were obtained from ECMWF on a 2.5° global grid. Although the analyses were permanently archived as spherical harmonic coefficients, ECMWF reconstituted the analyses for use in the data processing. Synoptic analyses at six-hour intervals were received for the six-year period, but only the 00 and 12Z analyses were re-sorted into a grid point sort. Given the quality control performed by ECMWF on collected data and the requirements for horizontal and vertical data consistency imposed by the assimilation system, minimal quality control was performed prior to summarization. Primary quality control was limited to comparison of level data against known/estimated climatological extremes.

The summarized grid point data were objectively analyzed, machine-contoured by parameter and level on polar stereographic (0°-90°N and S) and cylindrical equidistant (0°-60°N and S) projections with resulting contours machine-labeled. In addition, individual wind observations were consolidated into eight 45° segments centered on directions north, northeast, ... through northwest for display as wind roses on a series of cylindrical equidistant projections.

Since the ECMWF analyses were archived as spectral harmonic coefficients, the grid point reconstitution process provides data for all global 2.5° grid points. This naturally includes (for the 1000 through 700 mb levels) selected grid points at which the land elevations exceed the height of the pressure surface. For these grid points, a blanking program was used to eliminate both contours and grid point wind roses.

ANALYSES

1. Pressure-Height

Grid point geopotential height values (in dekameters) are summarized by month for 13 levels from 1000 mb to 30 mb with solid and dashed contours of mean values presented on pressure height charts. Standard deviation of height is calculated from the individual daily values with contours presented on a separate chart series including the standard deviation of vector mean wind. Local points of highest and lowest pressure are designated with H's and L's on the analyzed charts. Not all pressure centers are enclosed by closed contours. Vector mean wind in 5-knot increments are calculated for selected grid points considered adequate to depict flow for the hemisphere with wind shaft orientation related to specific latitude/longitude lines. Vector mean winds less than 2.5 knots are depicted as a shaft with no barbs. Contours of mean geopotential height and vector mean wind barbs are presented for the northern/southern hemispheres on polar stereographic projection and for 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

2. Wind Roses

Wind roses for 10° grid points from 5° to 85° north and south are presented by month for all levels from 1000 mb to 30 mb. Each hemisphere is divided into three longitudinal zones: 60°W to 60°E, 60°E to 180°E, and 180°W to 60°W. Each rose presents:

a) Scalar mean speed

b) Percent frequency of occurrence from each of 8 cardinal point wind directions proportional to shaft length with dots on the shafts representing 5 percentile intervals.

c) Mean speed for each of the 8 cardinal wind directions rounded to the nearest 5 knots.

Roses for grid points on the 1000 mb through 700 mb level charts are blanked whenever the land elevation exceeds the mean geopotential height of the specified level.

3. Temperature

Grid point temperature data (in °C) are summarized by month for 13 levels from 1000 mb to 30 mb with solid and dashed contours of mean values presented on pressure height charts. Temperature standard deviation derived from the individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

4. Dew Point

Grid point moisture data were received as mixing ratios for the period through April 19, 1982 and as relative humidity thereafter for the 1000 through 300 mb levels. All moisture data were converted to dew point values. These are summarized by month with solid and dashed contours of mean values presented on pressure height charts. Dew point standard deviation derived from the individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

5. Density

Grid point density data were computed from the daily values of temperature and pressure from the equation of state in the form

$$\rho = \frac{P}{RT}$$

where ρ is the density, P is the pressure, T is the temperature, and R is the gas constant. Density was computed for moist air through 300 mb and for dry air from 250 mb to 30 mb. Density data (in Kg/m^3) are summarized by month for all 13 levels with solid and dashed contours of mean values presented on pressure height charts. Density standard deviation derived from individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

6. Standard Deviation of Height and Vector Mean Wind

Standard deviation of the height and vector mean wind data presented on the pressure height charts are presented on monthly charts for the 1000 through 30 mb levels. Height standard deviations (in dekameters) are presented as solid contours and vector wind standard deviations (in knots) as dashed contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

7. Jet Stream

Grid point scalar mean wind speed (in knots), as presented by the value in the center of the wind rose octagons, are summarized by month and analyzed for 500 through 30 mb. All speeds exceeding 50 knots are shaded with shading intensity increasing by 25-knot increments. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections.

DATA AVAILABILITY

Monthly summarized grid point data for the period of record for all levels from 1000 through 30 mb have been retained on magnetic tape. Data available, per level, include:

Number of observations
Mean zonal wind component and standard deviation
Mean meridional wind component and standard deviation
Vector mean wind and standard deviation
Mean temperature and standard deviation
Mean dew point (through 300 mb) and standard deviation
Mean geopotential height and standard deviation
Mean density and standard deviation
Mean scalar wind speed and percentage of observations for each designated direction

Similarly summarized data for each half-month of the 1980-85 period are also available on magnetic tape. Summaries can be provided on magnetic media or in listing form by the National Climatic Data Center.

REFERENCES

ECMWF, 1988: User guide to ECMWF products.

Lorenc, A.C., 1981: A global three-dimensional multivariate statistical interpretation scheme. Monthly Weather Review, **109**, 701-721.

Lonnberg, P., J. Pailleux, and A. Hollingsworth, 1986: The new analyses system. ECMWF Technical Memorandum No. 125.

Naval Weather Service Command, 1969: Climate of the Upper Air - Southern Hemisphere, VOL I, Temperature, Dewpoint and Heights at Selected Pressure Levels, NAVAIR 50-1C-55.

Naval Weather Service Command, 1970: Selected Level Heights, Temperatures and Dewpoints for the Northern Hemisphere, NAVAIR 50-1C-52.

Shaw, D.B., P. Lonnberg, and A. Hollingsworth, 1984: The 1984 revision of the ECMWF Analysis System. ECMWF Technical Memorandum, No. 92.

U.S. Navy, 1959: Upper Wind Statistics Charts of the Northern Hemisphere, VOL I-III, NAVAIR 50-1C-535.

U.S. Navy, 1966: Components of the 1000 mb Winds of the Northern Hemisphere, NAVAIR 50-1C-51.

PRESSURE - HEIGHT
(13 LEVELS, 1000 TO 30 MB)

- Contours of mean height (solid and dashed lines) in geopotential dekameters; example: 580 is 5800 geopotential meters; solids labeled, dashed intermediates unlabeled

- Height labeled interval:

  6 dekameters (60 meters) - 1000 MB to 400 MB
  12 dekameters (120 meters) - 300 MB to 200 MB
  8 dekameters (80 meters) - 150 MB to 30 MB

- [illegible] for mean wind in knots

- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE



Mean Geopotential Height (dkm)

Upper Air Climatology

Vector Mean Wind (kt)

Northern Hemisphere

July

1981 MO





Upper Air Climatology

Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

1000 MB




Mean Geopotential Height (dkm)
Vector Mean Wind (kt)
Upper Air Climatology
Northern Hemisphere
156
150
144
156
150
156
156
156
156
156
156
156
150
90° W
0°
90° E




Mean Geopotential Height (dkm)
Vector Mean Wind (kt)
July
Upper Air Climatology
Northern Hemisphere
90° W
0°
90° E
180° E

Upper Air Climatology

Southern Hemisphere

Mean Geopotential Height (dam)

Vector Mean Wind (m/s)





Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

500 Mb



Upper Air Climatology
Southern Hemisphere

Mean Geopotential Height (dm)
Vector Mean Wind (kt)







Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

700 MB




Upper Air Climatology
Northern Hemisphere
Mean Pressure Height (dam)
Vector Mean Wind (m)
90°
L
950
940
924
900
876
852
828
804
780
756
732
708
684
660
636
612
588
564
540
516
492
468
444
420
396
372
348
324
300
276
252
228
204
180
156
132
108
84
60
36
12
-12
-36
-60
-84
-108
-132
-156
-180
-204
-228
-252
-276
-300
-324
-348
-372
-396
-420
-444
-468
-492
-516
-540
-564
-588
-612
-636
-660
-684
-708
-732
-756
-780
-804
-828
-852
-876
-900
-924
-948




Upper Air Climatology
Southern Hemisphere
Mean Geopotential Height (dam)
Vector Mean Wind (m/s)
July
1992
1988
1984
1980
1976
1972
1968
1964
1960
1956
1952
1948
1944
1940
1936
1932
1928
1924
1920
1916
1912
1908
1904
1900
1896
1892
1888
1884
1880
1876
1872
1868
1864
1860
1856
1852
1848
1844
1840
1836
1832
1828
1824
1820
1816
1812
1808
1804
1800
1796
1792
1788
1784
1780
1776
1772
1768
1764
1760
1756
1752
1748
1744
1740
1736
1732
1728
1724
1720
1716
1712
1708
1704
1700
1696
1692
1688
1684
1680
1676
1672
1668
1664
1660
1656
1652
1648
1644
1640
1636
1632
1628
1624
1620
1616
1612
1608
1604
1600
1596
1592
1588
1584
1580
1576
1572
1568
1564
1560
1556
1552
1548
1544
1540
1536
1532
1528
1524
1520
1516
1512
1508
1504
1500
1496
1492
1488
1484
1480
1476
1472
1468
1464
1460
1456
1452
1448
1444
1440
1436
1432
1428
1424
1420
1416
1412
1408
1404
1400
1396
1392
1388
1384
1380
1376
1372
1368
1364
1360
1356
1352
1348
1344
1340
1336
1332
1328
1324
1320
1316
1312
1308
1304
1300
1296
1292
1288
1284
1280
1276
1272
1268
1264
1260
1256
1252
1248
1244
1240
1236
1232
1228
1224
1220
1216
1212
1208
1204
1200
1196
1192
1188
1184
1180
1176
1172
1168
1164
1160
1156
1152
1148
1144
1140
1136
1132
1128
1124
1120
1116
1112
1108
1104
1100
1096
1092
1088
1084
1080
1076
1072
1068
1064
1060
1056
1052
1048
1044
1040
1036
1032
1028
1024
1020
1016
1012
1008
1004
1000
996
992
988
984
980
976
972
968
964
960
956
952
948
944
940
936
932
928
924
920
916
912
908
904
900
896
892
888
884
880
876
872
868
864
860
856
852
848
844
840
836
832
828
824
820
816
812
808
804
800
796
792
788
784
780
776
772
768
764
760
756
752
748
744
740
736
732
728
724
720
716
712
708
704
700
696
692
688
684
680
676
672
668
664
660
656
652
648
644
640
636
632
628
624
620
616
612
608
604
600
596
592
588
584
580
576
572
568
564
560
556
552
548
544
540
536
532
528
524
520
516
512
508
504
500
496
492
488
484
480
476
472
468
464
460
456
452
448
444
440
436
432
428
424
420
416
412
408
404
400
396
392
388
384
380
376
372
368
364
360
356
352
348
344
340
336
332
328
324
320
316
312
308
304
300
296
292
288
284
280
276
272
268
264
260
256
252
248
244
240
236
232
228
224
220
216
212
208
204
200
196
192
188
184
180
176
172
168
164
160
156
152
148
144
140
136
132
128
124
120
116
112
108
104
100
96
92
88
84
80
76
72
68
64
60
56
52
48
44
40
36
32
28
24
20
16
12
8
4
0
90°
60°
30°
0°
30° W
60° W
90° W
120° W
150° W
180°
150° E
120° E
90° E
60° E
30° E
0°
30° S
60° S
90° S

Mean Barometric Height (dkm)

Vector Mean Wind (kt)

July

500 MB

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology
Southern Hemisphere
Mean Geopotential Height
Vector Mean Wind
July







Upper Air Climatology

Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

100 MB



Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

1 MB

Upper Air Climatology

Northern Hemisphere





Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

July

50 MB









WIND ROSES
(13 LEVELS, 1000 TO 30 MB)

- Wind roses at 10 degree latitude/longitude grid points
- Directional mean wind speed in 5 knot increments
- Frequency proportional to barb length with individual dots representing 5% increments. Values greater than 30% are plotted directly on the barb.
- Roses blanked at grid points with elevations exceeding specified geopotential heights.
- Sample rose explanation:



ELEVATION SCALE







July

Upper Air Climatology
Southern Hemisphere





Italy
1000 MB

Upper Air Climatology
Northern Hemisphere














90° N
80° N
70° N
60° N
50° N
40° N
30° N
20° N







July
700 MB

Upper Air Climatology
Southern Hemisphere





Type A Climatology
Southern Hemisphere





July
500 Mb

Upper Air Climatology
Southern Hemisphere





July
500 MB

Upper Air Climatology
Southern Hemisphere





July
500 Mb

Upper Air Climatology
Southern Hemisphere





July
400 Mb

Upper Air Climatology
Southern Hemisphere



Upper Air Climatology
Southern Hemisphere





Upper Air Climatology
Southern Hemisphere





Upper Air Climatology
Southern Hemisphere



















July
850 MB

Upper Air Climatology
Southern Hemisphere





Upper Air Climatology
Southern Hemisphere





Upper Air Climatology
Southern Hemisphere





July

Upper Air Climatology
Southern Hemisphere




13 13 14 14 36 15 35 15 14 14 15 13 13 11
49 18 57 19 54 22 52 21 49 20 46 23 50 27 57 27 65 28 66 29 67 27 67 30
76 47 75 49 72 49 69 45 65 64 64 40 60 37 68 38 72 40 71 40 67 42 75 43
80 54 68 53 65 51 72 51 66 50 67 49 65 47 66 47 68 46 64 45 67 45 67 48
77 45 76 46 77 48 80 51 75 53 72 55 74 56 77 56 72 55 77 55 74 55 80 56
86 65 87 66 83 66 83 68 81 69 83 68 86 67 84 67 80 66 83 67 78 67 70 69
82 78 79 76 77 74 74 72 74 73 74 70 77 69 77 68 75 67 74 67 74 67 67 66
53 39 53 39 52 39 50 39 53 38 55 39 53 39 54 40 55 41 55 41 53 40 47 39 32
24 24 24 32 25 37 25 38 25 40 26 39 27 32 27 28 31 28 36 27 37

89


90° N
80° N
70° N
60° N
50° N
40° N
30° N
20° N
10° N
0°
60° W
50° W
40° E
50° E
60° E







July
70 MB

Upper Air Climatology
Southern Hemisphere





July
70 Mb

Upper Air Climatology
Southern Hemisphere













July

Upper Air Climatology
Southern Hemisphere









July
30 Mb

Upper Air Climatology
Southern Hemisphere





July

Upper Air Climatology
Southern Hemisphere

JET STREAM
(10 LEVELS, 500 TO 30 MB)

- Contours of mean scalar wind speed in knots
- Minimum mean scalar speed: 50 knots
- Contour interval of mean scalar speed: 25 knots

ELEVATION SCALE




Wind < 50 KT




Wind < 50 KT




50
50
50
50


Upper Air Climatology
Southern Hemisphere
50
75
50
75
50
75
75
50
75
50
75
50
75
75
75
50
50
50
50








50
50
50
50
50
50


50
50
50
50
50


50
75
100

50
50
50


50
75
50
90° S
50
50
75
50


50
75
50
75
50




50
75
50
90° S
75
50
50
50
50
75
50
75
50
180° W
90° W
0°
90° E
180° E



Upper Air Climatology

Southern Hemisphere

Jet Stream

50kt + 25kt Inc

July

70 MB








50
75
100
75
90° E
50
75
100
50
75
100
75
100
50
100
75

Jet Stream
50kt + 25kt inc
July
30 Mb



Upper Air Climatology

Southern Hemisphere

Jet Stream

50kt + 15kt inc

July

[illegible]



TEMPERATURE
(13 LEVELS, 1000 TO 30 MB)

- Contours of mean temperature (solid and dashed lines) in °C; solids labeled, dashed intermediates unlabeled
- Temperature labeled interval: 5°C
- Contours of standard deviation of temperature (dotted lines) in °C
- Standard deviation of temperature labeled interval: 2.5°C
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE






Std. Dev. < 2.5


90°
3.0
2.5

2.5
2.5
Std. Dev. < 2.5
2.5
Std. Dev. < 2.5
10




Std. Dev. < 2.5
Std. Dev. < 2.5


5.0
5.0
90°
180° W
90° W
0°
90° E



Std. Dev. < 2.5



Std. Dev. < 2.5


Std. Dev < 2.5
Std. Dev < 2.5


Std. Dev. < 2.5


Std. Dev. < 2.5
90° W
0°
90° E
180° E



Std. Dev. < 2.5



Std. Dev. < 2.5


5.0
Std. Dev. < 2.5



Std. Dev < 2.5



Std. Dev. < 2.5


Std. Dev. < 2.5


-30
Std. Dev. < 2.5





Std. Dev. < 2.5



Std. Dev. < 2.5




Std. Dev. < 2.5
Std. Dev. < 2.5


Upper Air Climatology
Southern Hemisphere
Mean Temperature (c)
Std Dev (Dotted)
July
180° W
90° W
0°
90° E
180° E

Mean Temperature (c)

Std Dev <Dotted>

July

70 Mb

Upper Air Climatology

Northern Hemisphere



Std. Dev. < 2.5



Std. Dev. < 2.5

Upper Air Climatology
Southern Hemisphere

Mean Temperature (c)
Std Dev <Dotted>
July
70 MB



Mean Temperature (c)

Std. Dev <Dotted>

July

50 Mb



Upper Air Climatology

Southern Hemisphere

Mean Temperature (c)

Std Dev <Dotted>

July

50 Mb



Mean Temperature (c)

Std Dev <Dotted>

July

10 Mb




180° W
90° W
0°
90° E
60° S

DEW POINT
(6 LEVELS, 1000 TO 300 MB)

- Contours of mean dew point (solid and dashed lines) in °C; solids labeled, dashed intermediates unlabeled.
- Dew point labeled interval: 5°C
- Contours of standard deviation of dew point (dotted lines) in °C
- Standard deviation of dew point labeled interval: 2.5°C
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE




Upper Air Climatology
Northern Hemisphere
90° W
0°
90° E
180° E






10.0
7.5
12.5
2.5
15
5.0
10
12.5
10.0


Upper Air Climatology
Northern Hemisphere
7.5
5.0
2.5
0
90° W
0°
90° E
180° E















DENSITY
(13 LEVELS, 1000 TO 30 MB)

- Contours of mean density (solid and dashed lines) in kilograms/cubic meter: solids labeled, dashed intermediates unlabeled

- Density labeled interval:

  .02 kilograms/cubic meter - 1000 MB to 400 MB
  .01 kilograms/cubic meter - 300 MB to 200 MB
  .006 kilograms/cubic meter - 150 MB to 30 MB

- Contours of standard deviation of density (dotted lines) in kilograms/cubic meter

- Standard deviation of density labeled interval:

  .01 kilograms/cubic meter - 1000 MB to 400 MB
  .005 kilograms/cubic meter - 300 MB to 200 MB
  .003 kilograms/cubic meter - 150 MB to 30 MB

- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE





Mean Density (kg/m³)

Std. Dev. (Dotted)

Upper Air Climatology

Northern Hemisphere




0.010
0.002
0.003
0.002
0.010
0.010
0.010
0.020


0.010
0.010
0.010
Std.Dev. < 0.010
0.010
0.840
Std.Dev. < 0.010





Std.Dev. < 0.010



Std.Dev. < 0.010


Std.Dev. < 0.010
Std.Dev. < 0.010
0.010
0.010



Std.Dev. < 0.010



Std.Dev. < 0.010



Std.Dev. < 0.010



Std.Dev. < 0.010



Std.Dev. < 0.005



Std.Dev. < 0.005



Std.Dev. < 0.005




Mean Density (kg/m3)
Std Dev (Dotted)
Upper Air Climatology
Northern Hemisphere
Std.Dev. < 0.005


Std.Dev. < 0.005


0.380
0.390
0.395
0.400
0.405
0.410
0.415
0.420
0.425
0.430
0.435
0.440
0.445
0.450
0.455
0.460
0.465
0.470
0.475
0.480
0.485
0.490
0.495
0.500
0.505
0.510
0.515
0.520
0.525
0.530
0.535
0.540
0.545
0.550
0.555
0.560
0.565
0.570
0.575
0.580
0.585


Std. Dev. < 0.105


0.005
0.320
Std. Dev. < 0.005


0.018
0.350
Std.Dev. < 0.005
Std.Dev. < 0.005

Mean Density (kg/m3)

Std Dev <Dotted>

July

150 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology
Southern Hemisphere

Mean Density (kg/m3)
Std Dev <Dotted>
July
150 Mb



Mean Density (kg/m3)

Std Dev <Dotted>

July

100 MB

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology
Southern Hemisphere
Mean Density (kg/m3)
Std Dev <Dotted>
July
100 MB
Std.Dev. < 0.003
Std.Dev. < 0.003



Std.Dev. < 0.003



Std.Dev. < 0.003

Upper Air Climatology
Southern Hemisphere

Mean Density (kg/m3)
Std Dev <Detrend>
July
71 Mb



Std.Dev. < 0.003



Mean Density (kg/m3)

Std Dev (Dotted)

July

90 KM

Upper Air Climatology

Northern Hemisphere






Upper Air Climatology
Southern Hemisphere
Mean Density (kg/m3)
Std Dev <Dotted>
July
50 Mb
0.084
0.092
90° B
Std.Dev. < 0.003
Std.Dev. < 0.003
0.084


Mean Density (kg/m3)
Std.Dev <Dotted>
July
30 MJD
Upper Air Climatology
Northern Hemisphere
0.045
Std.Dev. < 0.003
Density > 0.045
Density < 0.045
Std.Dev. > 0.003


Upper Air Climatology
Southern Hemisphere
Mean Density (kg/m3)
Std Dev (Dotted)
July
0.048
0.054
Std.Dev. < 0.003
0.048
0.048
0.048
0.048
0.048
Std.Dev. < 0.003

STANDARD DEVIATION OF HEIGHT
STANDARD DEVIATION OF VECTOR MEAN WIND
(13 LEVELS, 1000 TO 30 MB)

- Contours of standard deviation of height (solid lines) in geopotential dekameters

- Standard deviation of height labeled interval:

  3 dekameters (30 meters) - 1000 MB to 400 MB
  6 dekameters (60 meters) - 300 MB to 200 MB
  4 dekameters (40 meters) - 150 MB to 30 MB

- Contours of standard deviation of vector mean wind (dashed lines) in knots

- Standard deviation of vector mean wind labeled interval: 5 knots

- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE





10.0
9
10.0
9
5.0
5.0
10.0
9


10.0
5.0
5.0
5.0
5.0
5.0


10.0
10.0
10.0
10.0
5.0


10.0
5.0






25.0
12
9
10
20.0
5.0
5.0
5.0
15
15


20.0
5.0
20.0
5.0
20.0
5.0
5.0


25.0
15.0
12
10.0
20.0



Height (dkm) Std Dev <Solid>
Vector Std Dev (kt)
July
250 MB












10.0
5.0
5.0
10.0
10.0
10

Upper Air Climatology
Southern Hemisphere

Height (dkm) Std Dev <Solid>
Vector Std Dev (kt)
July
100 MB



10.0
10.0
10.0

10.0
15.0
20.0
25.0
12
24

Height (dkm) Std Dev <Solid>
Vector Std Dev (kt)
July
50 Mb






Height Std. Dev. < 12 dkm
Height Std. Dev. < 12 dkm